

On unification of the strong convergence theorems for a finite family of total asymptotically nonexpansive mappings in Banach spaces

FARRUKH MUKHAMEDOV

*Department of Computational & Theoretical Sciences
Faculty of Sciences, International Islamic University Malaysia
P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
e-mail: far75m@yandex.ru, farrukh_m@iiu.edu.my*

MANSOOR SABUROV

*Department of Computational & Theoretical Sciences
Faculty of Science, International Islamic University Malaysia
P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
e-mail: msaburov@gmail.com*

Abstract

In this paper, we unify all know iterative methods by introducing a new explicit iterative scheme for approximation of common fixed points of finite families of total asymptotically I -nonexpansive mappings. Note that such a scheme contains as a particular case of the method introduced in [C.E. Chidume, E.U. Ofoedu, *Inter. J. Math. & Math. Sci.* **2009**(2009) Article ID 615107, 17p]. We construct examples of total asymptotically nonexpansive mappings which are not asymptotically nonexpansive. Note that no such kind of examples were known in the literature. We prove the strong convergence theorems for such iterative process to a common fixed point of the finite family of total asymptotically I -nonexpansive and total asymptotically nonexpansive mappings, defined on a nonempty closed convex subset of uniformly convex Banach spaces. Moreover, our results extend and unify all known results.

Mathematics Subject Classification: 46N10; 46B20; 47H09; 47H10

Key words: Explicit iteration process; a total asymptotically I -nonexpansive mapping; a total asymptotically nonexpansive mapping; common fixed point.

1 Introduction

Let K be a nonempty subset of a real normed linear space X and $T : K \rightarrow K$ be a mapping. Denote by $F(T)$ the set of fixed points of T , that is, $F(T) = \{x \in K : Tx = x\}$. Throughout this paper, we always assume that X is a real Banach space and $F(T) \neq \emptyset$. Now let us recall some known definitions

Definition 1.1. A mapping $T : K \rightarrow K$ is said to be:

- (i) *nonexpansive*, if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in K$;
- (ii) *asymptotically nonexpansive*, if there exists a sequence $\{\lambda_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} \lambda_n = 1$ such that $\|T^n x - T^n y\| \leq \lambda_n \|x - y\|$ for all $x, y \in K$ and $n \in \mathbb{N}$;
- (iii) *asymptotically nonexpansive in the intermediate sense*, if it is continuous and the following inequality holds

$$\limsup_{n \rightarrow \infty} \sup_{x, y \in K} (\|T^n x - T^n y\| - \|x - y\|) \leq 0. \quad (1.1)$$

Remark 1.2. Observe that if we define

$$a_n := \sup_{x, y \in K} (\|T^n x - T^n y\| - \|x - y\|), \quad \sigma_n := \max\{0, a_n\}, \quad (1.2)$$

then $\sigma_n \rightarrow 0$ as $n \rightarrow \infty$ and (1.1) reduces to

$$\|T^n x - T^n y\| \leq \|x - y\| + \sigma_n, \quad \forall x, y \in K, n \geq 1. \quad (1.3)$$

In [3]-[4] Browder studied the iterative construction for fixed points of nonexpansive mappings on closed and convex subsets of a Hilbert space. Note that for the past 30 years or so, the study of the iterative processes for the approximation of fixed points of nonexpansive mappings and fixed points of some of their generalizations have been flourishing areas of research for many mathematicians (see for more details [15],[9]).

The class of asymptotically nonexpansive mappings was introduced by Goebel and Kirk [16] as a generalization of the class of nonexpansive mappings. They proved that if K is a nonempty closed convex bounded subset of a uniformly convex real Banach space and T is an asymptotically nonexpansive self-mapping of K , then T has a fixed point.

The class of mappings which are asymptotically nonexpansive in the intermediate sense was introduced by Bruck et al. [6]. It is known [20] that if K is a nonempty closed convex bounded subset of a uniformly convex Banach space X and $T : K \rightarrow K$ is an asymptotically nonexpansive mapping in the intermediate sense, then T has a fixed point. It is worth mentioning that the class of mappings which are asymptotically nonexpansive in the intermediate sense contains properly the class of asymptotically nonexpansive mappings (see, e.g., [21])

The iterative approximation problems for nonexpansive mapping, asymptotically nonexpansive mapping and asymptotically nonexpansive mapping in the intermediate sense were studied extensively in [16], [20], [21], [6], [5], [22],[37], [26], [17], [31],[32], [33],[7],[8],[13],[14].

There are many different types of concepts which generalize a notion of nonexpansive mapping. One of such concepts is a total asymptotically nonexpansive mapping ([1]) and second one is an asymptotically I -nonexpansive mapping ([30]). Let us recall some notions.

Definition 1.3. Let K be a nonempty closed subset of a real normed linear space X . $T : K \rightarrow K$ is called a total asymptotically nonexpansive mapping if there exist nonnegative real sequence $\{\mu_n\}$ and $\{\lambda_n\}$ with $\mu_n, \lambda_n \rightarrow 0$ as $n \rightarrow \infty$ and strictly increasing continuous function $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $\phi(0) = 0$ such that for all $x, y \in K$,

$$\|T^n x - T^n y\| \leq \|x - y\| + \mu_n \phi(\|x - y\|) + \lambda_n, \quad n \geq 1. \quad (1.4)$$

Remark 1.4. If $\phi(\xi) = \xi$, then (1.4) reduces to

$$\|T^n x - T^n y\| \leq (1 + \mu_n)\|x - y\| + \lambda_n, \quad n \geq 1. \quad (1.5)$$

In addition, if $\lambda_n = 0$ for all $n \geq 1$, then total asymptotically nonexpansive mappings coincide with asymptotically nonexpansive mappings. If $\mu_n = 0$ and $\lambda_n = 0$ for all $n \geq 1$, we obtain from (1.5) the class of mappings that includes the class of nonexpansive mappings. If $\mu_n = 0$ and $\lambda_n = \sigma_n = \max\{0, a_n\}$, where $a_n := \sup_{x, y \in K} (\|T^n x - T^n y\| - \|x - y\|)$ for all $n \geq 1$, then (1.5) reduces to (1.3) which has been studied as mappings asymptotically nonexpansive in the intermediate sense.

The idea of the definition of a total asymptotically nonexpansive mappings is that to unify various definitions of classes of mappings associated with the class of asymptotically nonexpansive mappings and to prove a general convergence theorems applicable to all these classes of nonlinear mappings.

Alber et.al. [1] studied methods of approximation of fixed points of total asymptotically nonexpansive mappings. Chidume et.al. [10] introduced an iterative scheme for approximation of a common fixed point of a finite family of total asymptotically nonexpansive mappings in Banach spaces. Recently, Chidume et.al. [11] constructed a new iterative sequence much simpler than other types of approximation of common fixed points of finite families of total asymptotically nonexpansive mappings.

On the other hand, in [30] an asymptotically I -nonexpansive mapping was introduced.

Definition 1.5. Let $T : K \rightarrow K$, $I : K \rightarrow K$ be two mappings of a nonempty subset K of a real normed linear space X . Then T is said to be:

- (i) I -nonexpansive, if $\|Tx - Ty\| \leq \|Ix - Iy\|$ for all $x, y \in K$;
- (ii) asymptotically I -nonexpansive, if there exists a sequence $\{\lambda_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} \lambda_n = 1$ such that $\|T^n x - T^n y\| \leq \lambda_n \|I^n x - I^n y\|$ for all $x, y \in K$ and $n \geq 1$;

Best approximation properties of I -nonexpansive mappings were investigated in [30, 12]. In [27] strong convergence of Mann iterations of I -nonexpansive mapping has been proved. In [23] the weak convergence of three-step Noor iterative scheme for an I -nonexpansive mapping in a Banach space has been established. In [34] the weakly convergence theorem for asymptotically I -nonexpansive mapping defined in Hilbert space was proved. Recently, in [18, 35, 36] the weak

and strong convergence of explicit and implicit iteration process to a common fixed point of a finite family of asymptotically I -nonexpansive mappings have been studied.

In this paper, we introduce a new type of concept of a generalization of nonexpansive mapping's notion, which is a combination of Definitions 1.3 and 1.5.

Definition 1.6. Let $T : K \rightarrow K$, $I : K \rightarrow K$ be two mappings of a nonempty subset K of a real normed linear space X . Then T is said to be a total asymptotically I -nonexpansive mapping if there exist nonnegative real sequences $\{\mu_n\}$ and $\{\lambda_n\}$ with $\mu_n, \lambda_n \rightarrow 0$ as $n \rightarrow \infty$ and the strictly increasing continuous function $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $\phi(0) = 0$ such that for all $x, y \in K$,

$$\|T^n x - T^n y\| \leq \|I^n x - I^n y\| + \mu_n \phi(\|I^n x - I^n y\|) + \lambda_n, \quad n \geq 1. \quad (1.6)$$

Now let us provide an examples of a total asymptotically I -nonexpansive mapping, which is not asymptotically nonexpansive mapping.

Example 1. Let us consider the space ℓ_1 , and let $B_1 = \{\mathbf{x} \in \ell_1 : \|\mathbf{x}\|_1 \leq 1\}$. Define a nonlinear operator $T : \ell_1 \rightarrow \ell_1$ by

$$T_\alpha(x_1, x_2, \dots, x_n, \dots) = (0, \alpha\sqrt{|x_1|}, \alpha x_2, \dots, \alpha x_n, \dots), \quad \alpha \in (0, 1). \quad (1.7)$$

Let $\|\mathbf{x}\|_1 \leq 1$, then from

$$\begin{aligned} \|T_\alpha(\mathbf{x})\|_1 &= \alpha \left(\|\mathbf{x}\|_1 - |x_1| + \sqrt{|x_1|} \right) \\ &\leq \alpha \left(1 - |x_1| + \sqrt{|x_1|} \right) \leq 1 \end{aligned}$$

one gets $T(B_1) \subset B_1$.

One can find that

$$T_\alpha^k(x_1, x_2, \dots, x_n, \dots) = (\underbrace{0, \dots, 0}_k, \alpha^k \sqrt{|x_1|}, \alpha^k x_2, \dots, \alpha^k x_n, \dots).$$

Hence,

$$\|T_\alpha^k(\mathbf{x}) - T_\alpha^k(\mathbf{y})\|_1 = \alpha^k \left(\|\mathbf{x} - \mathbf{y}\|_1 + |\sqrt{|x_1|} - \sqrt{|y_1|}| - |x_1 - y_1| \right). \quad (1.8)$$

From $\mathbf{x}, \mathbf{y} \in B_1$ we have

$$|\sqrt{|x_1|} - \sqrt{|y_1|}| \leq \sqrt{||x_1| - |y_1||} \leq \sqrt{\|\mathbf{x} - \mathbf{y}\|_1}. \quad (1.9)$$

So, it follows from (1.8), (1.9) that

$$\|T_\alpha^k(\mathbf{x}) - T_\alpha^k(\mathbf{y})\|_1 \leq \alpha^k \left(\|\mathbf{x} - \mathbf{y}\|_1 + \sqrt{\|\mathbf{x} - \mathbf{y}\|_1} \right) \quad \text{for all } \mathbf{x}, \mathbf{y} \in B_1, \quad k \in \mathbb{N}. \quad (1.10)$$

Now consider a new Banach space $\mathbb{R} \times \ell_1$ with a norm $\|\mathbb{X}\| = |x| + \|\mathbf{x}\|_1$, where $\mathbb{X} = (x, \mathbf{x})$ and define a new mapping $S : \mathbb{R} \times \ell_1 \rightarrow \mathbb{R} \times \ell_1$ by

$$S(x, \mathbf{x}) = (x, T_\alpha(\mathbf{x})).$$

Let $K = [0, 1] \times B_1$. Then it is clear that $S(K) \subset K$. One can see that $S^k(x, \mathbf{x}) = (x, T_\alpha^k(\mathbf{x}))$. Therefore, using (1.11) we obtain

$$\begin{aligned} \|S^k(\mathbb{X}) - S^k(\mathbb{Y})\| &= |x - y| + \|T_\alpha^k(\mathbf{x}) - T_\alpha^k(\mathbf{y})\|_1 \\ &\leq |x - y| + \alpha^k \left(\|\mathbf{x} - \mathbf{y}\|_1 + \sqrt{\|\mathbf{x} - \mathbf{y}\|_1} \right) \\ &\leq \|\mathbb{X} - \mathbb{Y}\| + \alpha^k \left(\|\mathbb{X} - \mathbb{Y}\| + \sqrt{\|\mathbb{X} - \mathbb{Y}\|} \right) \end{aligned} \quad (1.11)$$

We let $\phi(t) = t + \sqrt{t}$ and $\mu_k = \alpha^k$. It is clear that $\phi(0) = 0$ and ϕ is strictly increasing, and moreover (1.11) implies

$$\|S^k(\mathbb{X}) - S^k(\mathbb{Y})\| \leq \|I\mathbb{X} - I\mathbb{Y}\| + \mu_k \phi(\|I\mathbb{X} - I\mathbb{Y}\|)$$

that S is a totally asymptotically I -nonexpansive mapping. Here I is the identity mapping of $\mathbb{R} \times \ell_1$.

Now we are going to show that S is not asymptotically nonexpansive. Namely, we will establish that for any sequence of positive numbers $\{\lambda_n\}$ with $\lambda_n \rightarrow 0$ and any $k \in \mathbb{N}$ one can find $\mathbb{X}_0, \mathbb{Y}_0$ such that

$$\|S^k(\mathbb{X}_0) - S^k(\mathbb{Y}_0)\| > (1 + \lambda_k) \|\mathbb{X}_0 - \mathbb{Y}_0\|.$$

In fact, choose $\mathbb{X}_0, \mathbb{Y}_0$ as follows:

$$\mathbb{X}_0 = (0, \mathbf{x}_0), \quad \mathbb{Y}_0 = (0, \mathbf{y}_0),$$

where

$$\mathbf{x}_0 = (x_0, 0, \dots, 0, \dots), \quad \mathbf{y}_0 = \left(\frac{x_0}{4}, 0, \dots, 0, \dots\right)$$

and

$$0 < x_0 < \frac{4\alpha^{2k}}{9(1 + \lambda_k)^2}. \quad (1.12)$$

From (1.8) one finds

$$\begin{aligned} \|S^k(\mathbb{X}_0) - S^k(\mathbb{Y}_0)\| &= \alpha^k \left| \sqrt{x_0} - \frac{\sqrt{x_0}}{2} \right| = \alpha^k \frac{\sqrt{x_0}}{2}, \\ \|\mathbb{X}_0 - \mathbb{Y}_0\| &= \frac{3x_0}{4}. \end{aligned}$$

The last equalities with (1.12) imply that

$$\frac{\|S^k(\mathbb{X}_0) - S^k(\mathbb{Y}_0)\|}{\|\mathbb{X}_0 - \mathbb{Y}_0\|} = \frac{2\alpha^k}{3\sqrt{x_0}} > 1 + \lambda_k.$$

This yields the required assertion. Note that S has infinitely many fixed points in K , i.e. $\text{Fix}(S) = \{(x, \mathbf{0}) : x \in [0, 1]\}$.

EXAMPLE 2. Let us consider the Banach space $\mathbb{R} \times \ell_1$ defined as before, and f be a mapping of a segment $C \subset \mathbb{R}$ to itself, i.e. $f : C \rightarrow C$ with $f(0) = 0$ and

$$|f^n(x) - f^n(y)| \leq |x - y| + c_n, \quad c_n > 0, \quad n \in \mathbb{N}$$

where $c_n \rightarrow 0$. Note that such kind of functions do exist. One can take (see for more details [21]) $C = [-1/\pi, 1/\pi]$ and

$$f_\kappa(x) = \begin{cases} \kappa x \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0. \end{cases} \quad \kappa \in (0, 1).$$

Define a new mapping $S_f : C \times B_1 \rightarrow C \times B_1$ by

$$S_f(x, \mathbf{x}) = (f(x), T_\alpha(\mathbf{x})),$$

here T is defined as above (see (1.7)). Using the same argument as above Example 1, we can establish that

$$\|S_f^k(\mathbb{X}) - S_f^k(\mathbb{Y})\| \leq \|\mathbb{X} - \mathbb{Y}\| + \mu_k \phi(\|\mathbb{X} - \mathbb{Y}\|) + c_k \quad \forall k \in \mathbb{N}.$$

Moreover, such a mapping is not asymptotically nonexpansive. Note that the mapping S_{f_κ} with the function f_κ has a unique fixed point in $C \times B_1$.

Remark. To the best of our knowledge, we should stress that the constructed examples are currently only unique examples of totally asymptotically nonexpansive mappings which are not asymptotically nonexpansive. Before, no such examples were known in the literature.

Aim of the present paper is unification of all known iterative methods by introducing a new iterative scheme for approximation of common fixed points of finite families of total asymptotically I -nonexpansive mappings. Note that such a scheme contains as a particular case of the method introduced in [11], and allow us to construct more simpler methods than [10, 11].

Namely, let K be a nonempty closed convex subset of a real Banach space X and $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive mappings, i.e.

$$\|T_i^n x - T_i^n y\| \leq \|I_i^n x - I_i^n y\| + \mu_{in} \phi_i(\|I_i^n x - I_i^n y\|) + \lambda_{in} \quad (1.13)$$

and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically nonexpansive mappings, i.e.

$$\|I_i^n x - I_i^n y\| \leq \|x - y\| + \tilde{\mu}_{in} \varphi_i(\|x - y\|) + \tilde{\lambda}_{in}, \quad (1.14)$$

here $\phi_i, \varphi_i : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are the strictly increasing continuous functions with $\phi_i(0) = \varphi_i(0) = 0$ for all $i = \overline{1, m}$ and $\{\mu_{in}\}_{n=1}^\infty, \{\lambda_{in}\}_{n=1}^\infty, \{\tilde{\mu}_{in}\}_{n=1}^\infty, \{\tilde{\lambda}_{in}\}_{n=1}^\infty$ are nonnegative real sequences with

$\mu_{in}, \lambda_{in}, \tilde{\mu}_{in}, \tilde{\lambda}_{in} \rightarrow 0$ as $n \rightarrow \infty$ for all $i = \overline{1, m}$. Then for given sequences $\{\alpha_{jn}\}_{n=1}^{\infty}$, $\{\beta_{jn}\}_{n=1}^{\infty}$ in $(0, 1)$, where $j = \overline{0, m}$, we shall consider the following explicit iterative process:

$$\begin{cases} x_0 \in K, \\ x_{n+1} = \alpha_{0n}x_n + \sum_{i=1}^m \alpha_{in}T_i^n y_n \\ y_n = \beta_{0n}x_n + \sum_{i=1}^m \beta_{in}I_i^n x_n. \end{cases} \quad (1.15)$$

such that $\sum_{j=0}^m \alpha_{jn} = 1$ and $\sum_{j=0}^m \beta_{jn} = 1$.

Chidume et.al. [11] has considered only a particular case of the explicit iterative process (1.15), in which $\{I_i\}_{i=1}^m$ to be taken as the identity mappings. One of the main results of [11] (see Theorem 3.5. p.11) was correct while the provided proof of that result was wrong. Since, in their proof they used Lemma 2.3, but which actually is not applicable in that situation, since the sequence $\{t_n\}_{n=1}^{\infty}$ tends to 0. As a counterexample, we can consider the following one: let $x \in X$, $\|x\| = d > 0$, and let the sequences x_n, y_n, t_n be defined as follows

$$x_n = x, \quad y_n = -x, \quad t_n = \frac{1}{n}, \quad \forall n \in \mathbb{N}.$$

It is then clear that

$$\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n) y_n\| = \|x\| \lim_{n \rightarrow \infty} \left| 1 - \frac{2}{n} \right| = d.$$

However,

$$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 2d > 0.$$

In this paper, we shall provide a correct proof of Theorem 3.5 p.11 in [11]. As we already mentioned Lemma 2.3 is not applicable the main result of [11]. Therefore, we first will generalize Lemma 2.3 to the case of finite number of sequences. Such a generalization gives us a possibility to prove the mentioned result. On other hand, the provided generalization presents an independent interest as well. Moreover, we extend and unify the main result of [11] for a finite family of total asymptotically I_i -nonexpansive mappings $\{T_i\}_{i=1}^m$. Namely, we shall prove the strong convergence of the explicit iterative process (1.15) to a common fixed point of the finite family of total asymptotically I_i -nonexpansive mappings $\{T_i\}_{i=1}^m$ and the finite family of total asymptotically nonexpansive mappings $\{I_i\}_{i=1}^m$. Here, we stress that Lemmas 3.1 and 3.2 play a crucial role. All presented results here extend, generalize, unify and improve the corresponding main results of [1], [11], [18], [35],[36],[28], [29,].

2 Preliminaries

Throughout this paper, we always assume that X is a real Banach space. The following lemmas play an important role in proving our main results.

Lemma 2.1. [33] Let $\{a_n\}, \{b_n\}, \{c_n\}$ be three sequences of nonnegative real numbers with $\sum_{n=1}^{\infty} b_n < \infty, \sum_{n=1}^{\infty} c_n < \infty$. If the following condition is satisfied

$$(i) \quad a_{n+1} \leq (1 + b_n)a_n + c_n, \quad n \geq 1.$$

then the limit $\lim_{n \rightarrow \infty} a_n$ exists.

Lemma 2.2. [38] Let X be a uniformly convex Banach space and $t \in (0, 1)$. Suppose $\{x_n\}, \{y_n\}$ are two sequences in X such that

$$\lim_{n \rightarrow \infty} \|tx_n + (1 - t)y_n\| = d, \quad \limsup_{n \rightarrow \infty} \|x_n\| \leq d, \quad \limsup_{n \rightarrow \infty} \|y_n\| \leq d,$$

holds some $d \geq 0$. Then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.

Lemma 2.3. [31] Let X be a uniformly convex Banach space and b, c be two constants with $0 < b < c < 1$. Suppose that $\{t_n\}$ is a sequence in $[b, c]$ and $\{x_n\}, \{y_n\}$ are two sequences in X such that

$$\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n)y_n\| = d, \quad \limsup_{n \rightarrow \infty} \|x_n\| \leq d, \quad \limsup_{n \rightarrow \infty} \|y_n\| \leq d,$$

holds some $d \geq 0$. Then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.

3 Main results

In this section we shall prove our main results. To formulate ones, we need some auxiliary results.

First we are going to generalize of Lemmas 2.2 and 2.3 for m number of sequences $\{z_{in}\}_{n=1}^{\infty}$ from the uniformly convex Banach space X , where $i = \overline{1, m}$.

Lemma 3.1. Let X be a uniformly convex Banach space and $\alpha_i \in (0, 1), i = \overline{1, m}$ be any constants with $\sum_{i=1}^m \alpha_i = 1$. Suppose $\{z_{in}\}_{n=1}^{\infty}, i = \overline{1, m}$ are sequences in X such that

$$\lim_{n \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_i z_{in} \right\| = d, \quad \limsup_{n \rightarrow \infty} \|z_{in}\| \leq d, \quad \forall i = \overline{1, m}, \quad (3.1)$$

holds some $d \geq 0$. Then $\lim_{n \rightarrow \infty} \|z_{in}\| = d$ and $\lim_{n \rightarrow \infty} \|z_{in} - z_{jn}\| = 0$ for any $i, j = \overline{1, m}$.

Proof. Let us first prove $\lim_{n \rightarrow \infty} \|z_{in}\| = d$ for any $i = \overline{1, m}$. Indeed, it follows from (3.1) that

$$\begin{aligned} d &= \lim_{n \rightarrow \infty} \left\| \sum_{k=1}^m \alpha_k z_{kn} \right\| = \liminf_{n \rightarrow \infty} \left\| \sum_{k=1}^m \alpha_k z_{kn} \right\| \\ &\leq \liminf_{n \rightarrow \infty} \left(\sum_{k=1}^m \alpha_k \|z_{kn}\| \right) \leq \alpha_i \liminf_{n \rightarrow \infty} \|z_{in}\| + \sum_{k \neq i} \alpha_k \limsup_{n \rightarrow \infty} \|z_{kn}\| \\ &\leq \alpha_i \liminf_{n \rightarrow \infty} \|z_{in}\| + (1 - \alpha_i)d. \end{aligned}$$

We then get that $\liminf_{n \rightarrow \infty} \|z_{in}\| \geq d$, which means $\lim_{n \rightarrow \infty} \|z_{in}\| = d$.

Now we prove the statement $\lim_{n \rightarrow \infty} \|z_{in} - z_{jn}\| = 0$ by means of mathematical induction w.r.t. m . For $m = 2$, the statement immediately follows from Lemma 2.2. Assume that the statement is true, for $m = k - 1$. Let us prove for $m = k$. To do this, denote

$$t_n = \frac{1}{1 - \alpha_k} \sum_{i=1}^{k-1} \alpha_i z_{in}.$$

Since $\frac{1}{1 - \alpha_k} \sum_{i=1}^{k-1} \alpha_i = 1$ we get $\limsup_{n \rightarrow \infty} \|t_n\| \leq d$. On the other hand, one has

$$\begin{aligned} d &= \liminf_{n \rightarrow \infty} \left\| \sum_{i=1}^k \alpha_i z_{in} \right\| = \liminf_{n \rightarrow \infty} \|(1 - \alpha_k)t_n + \alpha_k z_{kn}\| \\ &\leq (1 - \alpha_k) \liminf_{n \rightarrow \infty} \|t_n\| + \alpha_k \limsup_{n \rightarrow \infty} \|z_{kn}\| \\ &\leq (1 - \alpha_k) \liminf_{n \rightarrow \infty} \|t_n\| + \alpha_k d. \end{aligned}$$

We then obtain $\liminf_{n \rightarrow \infty} \|t_n\| \geq d$ which means $\lim_{n \rightarrow \infty} \|t_n\| = d$. In this case, according to the assumption of induction with the sequence t_n , we can conclude that $\lim_{n \rightarrow \infty} \|z_{in} - z_{jn}\| = 0$, if $1 \leq i, j \leq k - 1$.

Since $\lim_{n \rightarrow \infty} \|(1 - \alpha_k)t_n + \alpha_k z_{kn}\| = d$ due to Lemma 2.2 one gets

$$\lim_{n \rightarrow \infty} \|t_n - z_{kn}\| = 0.$$

If $1 \leq j \leq k - 1$ then the following inequality

$$\begin{aligned} \|z_{jn} - z_{kn}\| &\leq \|z_{jn} - t_n\| + \|t_n - z_{kn}\| \\ &\leq \frac{1}{1 - \alpha_k} \sum_{i=1}^{k-1} \alpha_i \|z_{in} - z_{jn}\| + \|t_n - z_{kn}\| \end{aligned}$$

implies that $\lim_{n \rightarrow \infty} \|z_{jn} - z_{kn}\| = 0$. This completes the proof. \square

Lemma 3.2. *Let X be a uniformly convex Banach space and α_*, α^* be two constants with $0 < \alpha_* < \alpha^* < 1$. Suppose that $\{\alpha_{in}\}_{n=1}^\infty \subset [\alpha_*, \alpha^*]$, $i = \overline{1, m}$ are any sequences with $\sum_{i=1}^m \alpha_{in} = 1$ for all $n \in \mathbb{N}$. Suppose $\{z_{in}\}_{n=1}^\infty, i = \overline{1, m}$ are sequences in X such that*

$$\lim_{n \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_{in} z_{in} \right\| = d, \quad \limsup_{n \rightarrow \infty} \|z_{in}\| \leq d, \quad \forall i = \overline{1, m}, \quad (3.2)$$

holds for some $d \geq 0$. Then $\lim_{n \rightarrow \infty} \|z_{in}\| = d$ and $\lim_{n \rightarrow \infty} \|z_{in} - z_{jn}\| = 0$ for any $i, j = \overline{1, m}$.

Proof. Analogously as in the proof of Lemma 3.1, it is easy to show that $\lim_{n \rightarrow \infty} \|z_{in}\| = d$. Therefore, let us prove the statement $\lim_{n \rightarrow \infty} \|z_{in} - z_{jn}\| = 0$ for any $i, j = \overline{1, m}$. Suppose contrary, i.e., there exist two numbers i_0, j_0 such that

$$\limsup_{n \rightarrow \infty} \|z_{i_0 n} - z_{j_0 n}\| = \beta_{i_0, j_0} > 0.$$

Then, there exists a subsequence $\{z_{i_0 n_k} - z_{j_0 n_k}\}_{k=1}^\infty$ of $\{z_{i_0 n} - z_{j_0 n}\}_{n=1}^\infty$ such that $\lim_{k \rightarrow \infty} \|z_{i_0 n_k} - z_{j_0 n_k}\| = \beta_{i_0 j_0}$.

Let us consider the subsequences $\{\alpha_{in_k}\}_{k=1}^\infty$ of $\{\alpha_{in}\}_{n=1}^\infty$, here $i = \overline{1, m}$. Since $\{\alpha_{in_k}\}_{k=1}^\infty \subset [\alpha_*, \alpha^*]$ there exist a subsequence $\{n_{k_l}\}_{l=1}^\infty$ of $\{n_k\}_{k=1}^\infty$ such that $\lim_{l \rightarrow \infty} \alpha_{in_{k_l}} = \alpha_i$ for all $i = \overline{1, m}$. Since $\sum_{i=1}^m \alpha_{in} = 1$, for all $n \in \mathbb{N}$, one gets $\sum_{i=1}^m \alpha_i = 1$, and $\alpha_i \in [\alpha_*, \alpha^*]$, for all $i = \overline{1, m}$. We know that

$$\begin{aligned} d &= \lim_{l \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_{in_{k_l}} z_{in_{k_l}} \right\| = \liminf_{l \rightarrow \infty} \left\| \sum_{i=1}^m \left((\alpha_{in_{k_l}} - \alpha_i) z_{in_{k_l}} + \alpha_i z_{in_{k_l}} \right) \right\| \\ &\leq \liminf_{l \rightarrow \infty} \left(\sum_{i=1}^m |\alpha_{in_{k_l}} - \alpha_i| \|z_{in_{k_l}}\| + \left\| \sum_{i=1}^m \alpha_i z_{in_{k_l}} \right\| \right) \\ &\leq \sum_{i=1}^m \limsup_{l \rightarrow \infty} \left(|\alpha_{in_{k_l}} - \alpha_i| \|z_{in_{k_l}}\| \right) + \liminf_{l \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_i z_{in_{k_l}} \right\| \end{aligned}$$

It then follows that $\liminf_{l \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_i z_{in_{k_l}} \right\| \geq d$. On the other hand, we have

$$\limsup_{l \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_i z_{in_{k_l}} \right\| \leq \sum_{i=1}^m \alpha_i \limsup_{l \rightarrow \infty} \|z_{in_{k_l}}\| \leq d.$$

Therefore, $\lim_{l \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_i z_{in_{k_l}} \right\| = d$. Consequently, Lemma 3.1 implies that $\lim_{l \rightarrow \infty} \|z_{i_0 n_{k_l}} - z_{j_0 n_{k_l}}\| = 0$. However, it contradicts to

$$\lim_{k \rightarrow \infty} \|z_{i_0 n_k} - z_{j_0 n_k}\| = \lim_{k \rightarrow \infty} \|z_{i_0 n_k} - z_{j_0 n_k}\| = \beta_{i_0 j_0} > 0.$$

This completes the proof. \square

Proposition 3.3. *Let X be a real Banach space and K be a nonempty closed convex subset of X . Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive mappings with sequences $\{\mu_{in}\}_{n=1}^\infty, \{\lambda_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$, and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family total asymptotically nonexpansive mappings with sequences $\{\tilde{\mu}_{in}\}_{n=1}^\infty, \{\tilde{\lambda}_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$. Suppose that there exist $M_i, M_i^*, N_i, N_i^* > 0$, $i = \overline{1, m}$ such that $\phi_i(\xi_i) \leq M_i^* \xi_i$, for all $\xi_i \geq M_i$ and $\varphi_i(\zeta_i) \leq N_i^* \zeta_i$ for all $\zeta_i \geq N_i$, where $i = \overline{1, m}$. Then the following holds for any $x, y \in K$ and for any $i = \overline{1, m}$,*

$$\|I_i^n x - I_i^n y\| \leq (1 + \tilde{\mu}_{in} N_i^*) \|x - y\| + \tilde{\mu}_{in} \varphi_i(N_i) + \tilde{\lambda}_{in}, \quad (3.3)$$

$$\begin{aligned} \|T_i^n x - T_i^n y\| &\leq (1 + \mu_{in} M_i^*) (1 + \tilde{\mu}_{in} N_i^*) \|x - y\| + \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) \\ &\quad + \tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) + \mu_{in} \phi_i(M_i) + \lambda_{in}, \end{aligned} \quad (3.4)$$

Proof. Since $\phi_i, \varphi_i : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are the strictly increasing continuous functions, where $i = \overline{1, m}$, it follows that $\phi_i(\xi_i) \leq \phi_i(M_i)$ and $\varphi_i(\zeta_i) \leq \varphi_i(N_i)$ whenever $\xi_i \leq M_i$ and $\zeta_i \leq N_i$, where

$i = \overline{1, m}$. By hypothesis of Proposition 3.3, for all $\xi_i, \zeta_i \geq 0$ and $i = \overline{1, m}$, we then get

$$\phi_i(\xi_i) \leq \phi_i(M_i) + M_i^* \xi_i, \quad (3.5)$$

$$\varphi_i(\zeta_i) \leq \varphi_i(N_i) + N_i^* \zeta_i. \quad (3.6)$$

Since $\{T_i\}_{i=1}^m : K \rightarrow K$, $\{I_i\}_{i=1}^m : K \rightarrow K$ are a total asymptotically I_i -nonexpansive and a total asymptotically nonexpansive mappings, respectively, from (3.5) and (3.6) one gets

$$\begin{aligned} \|I_i^n x - I_i^n y\| &\leq \|x - y\| + \tilde{\mu}_{in} \varphi_i(\|x - y\|) + \tilde{\lambda}_{in} \\ &\leq \|x - y\| + \tilde{\mu}_{in} (\varphi_i(N_i) + N_i^* \|x - y\|) + \tilde{\lambda}_{in} \\ &= (1 + \tilde{\mu}_{in} N_i^*) \|x - y\| + \tilde{\mu}_{in} \varphi_i(N_i) + \tilde{\lambda}_{in} \end{aligned}$$

and

$$\begin{aligned} \|T_i^n x - T_i^n y\| &\leq \|I_i^n x - I_i^n y\| + \mu_{in} \phi_i(\|I_i^n x - I_i^n y\|) + \lambda_{in} \\ &\leq \|I_i^n x - I_i^n y\| + \mu_{in} (\phi_i(M_i) + M_i^* \|I_i^n x - I_i^n y\|) + \lambda_{in} \\ &= (1 + \mu_{in} M_i^*) \|I_i^n x - I_i^n y\| + \mu_{in} \phi_i(M_i) + \lambda_{in} \\ &\leq (1 + \mu_{in} M_i^*) (1 + \tilde{\mu}_{in} N_i^*) \|x - y\| + \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) \\ &\quad + \tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) + \mu_{in} \phi_i(M_i) + \lambda_{in} \end{aligned}$$

□

Lemma 3.4. *Let X be a uniformly convex real Banach space and K be a nonempty closed convex subset of X . Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive mappings with sequences $\{\mu_{in}\}_{n=1}^\infty, \{\lambda_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$, and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family total asymptotically nonexpansive mappings with sequences $\{\tilde{\mu}_{in}\}_{n=1}^\infty, \{\tilde{\lambda}_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$, such that $F := \bigcap_{i=1}^m (F(T_i) \cap F(I_i)) \neq \emptyset$. Suppose $\sum_{n=1}^\infty \mu_{in} < \infty$, $\sum_{n=1}^\infty \lambda_{in} < \infty$, $\sum_{n=1}^\infty \tilde{\mu}_{in} < \infty$, $\sum_{n=1}^\infty \tilde{\lambda}_{in} < \infty$ for all $i = \overline{1, m}$ and there exist $M_i, M_i^*, N_i, N_i^* > 0$, $i = \overline{1, m}$ such that $\phi_i(\xi_i) \leq M_i^* \xi_i$, for all $\xi_i \geq M_i$ and $\varphi_i(\zeta_i) \leq N_i^* \zeta_i$ for all $\zeta_i \geq N_i$, where $i = \overline{1, m}$. If $\{x_n\}$ is the explicit iterative sequence defined by (1.15) then for each $p \in F$ the limit $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists.*

Proof. Since $F \neq \emptyset$, for any given $p \in F$, it follows from (1.15) and (3.4) that

$$\begin{aligned}
\|x_{n+1} - p\| &= \left\| \left(1 - \sum_{i=1}^m \alpha_{in}\right) (x_n - p) + \sum_{i=1}^m \alpha_{in} (T_i^n y_n - p) \right\| \\
&\leq \left(1 - \sum_{i=1}^m \alpha_{in}\right) \|x_n - p\| + \sum_{i=1}^m \alpha_{in} \|T_i^n y_n - p\| \\
&\leq \left(1 - \sum_{i=1}^m \alpha_{in}\right) \|x_n - p\| \\
&\quad + \sum_{i=1}^m \alpha_{in} (1 + \mu_{in} M_i^*) (1 + \tilde{\mu}_{in} N_i^*) \|y_n - p\| \\
&\quad + \sum_{i=1}^m \left(\alpha_{in} \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) + \alpha_{in} \tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) \right) \\
&\quad + \sum_{i=1}^m (\alpha_{in} \mu_{in} \phi_i(M_i) + \alpha_{in} \lambda_{in}). \tag{3.7}
\end{aligned}$$

Again from (1.15) and (3.3) we derive that

$$\begin{aligned}
\|y_n - p\| &= \left\| \left(1 - \sum_{i=1}^m \beta_{in}\right) (x_n - p) + \sum_{i=1}^m \beta_{in} (I_i^n x_n - p) \right\| \\
&\leq \left(1 - \sum_{i=1}^m \beta_{in}\right) \|x_n - p\| + \sum_{i=1}^m \beta_{in} \|I_i^n x_n - p\| \\
&= \left(1 + \sum_{i=1}^m \tilde{\mu}_{in} \beta_{in} N_i^*\right) \|x_n - p\| \\
&\quad + \sum_{i=1}^m \left(\tilde{\mu}_{in} \beta_{in} \varphi_i(N_i) + \tilde{\lambda}_{in} \beta_{in} \right) \tag{3.8}
\end{aligned}$$

Then from (3.7) and (3.8) one finds

$$\|x_{n+1} - p\| \leq (1 + b_n) \|x_n - p\| + c_n \tag{3.9}$$

here

$$\begin{aligned}
b_n &= \sum_{i=1}^m \left(\mu_{in} \alpha_{in} M_i^* + \tilde{\mu}_{in} \alpha_{in} N_i^* + \alpha_{in} \sum_{i=1}^m \tilde{\mu}_{in} \beta_{in} N_i^* \right) \\
&\quad + \sum_{i=1}^m \mu_{in} \tilde{\mu}_{in} \alpha_{in} M_i^* N_i^* + \sum_{i=1}^m \mu_{in} \alpha_{in} M_i^* \cdot \sum_{i=1}^m \tilde{\mu}_{in} \beta_{in} N_i^* \\
&\quad + \sum_{i=1}^m \tilde{\mu}_{in} \alpha_{in} N_i^* \cdot \sum_{i=1}^m \tilde{\mu}_{in} \beta_{in} N_i^*, \\
c_n &= \sum_{i=1}^m (\tilde{\mu}_{in} \beta_{in} \varphi_i(N_i) + \tilde{\lambda}_{in} \beta_{in}) \cdot \sum_{i=1}^m \alpha_{in} (1 + \mu_{in} M_i^*) (1 + \tilde{\mu}_{in} N_i^*) \\
&\quad + \sum_{i=1}^m \left(\tilde{\mu}_{in} \alpha_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) + \tilde{\lambda}_{in} \alpha_{in} (1 + \mu_{in} M_i^*) \right) \\
&\quad + \sum_{i=1}^m (\mu_{in} \alpha_{in} \phi_i(M_i) + \lambda_{in} \alpha_{in})
\end{aligned}$$

Denoting $a_n = \|x_n - p\|$ in (3.9) one gets

$$a_{n+1} \leq (1 + b_n) a_n + c_n.$$

Since $\sum_{n=1}^{\infty} b_n < \infty$ and $\sum_{n=1}^{\infty} c_n < \infty$, it follows from Lemma 2.1 the existence of the limit $\lim_{n \rightarrow \infty} a_n$. This means the limit

$$\lim_{n \rightarrow \infty} \|x_n - p\| = d \quad (3.10)$$

exists, where $d \geq 0$ is a constant. This completes the proof. \square

Now we prove the following result.

Theorem 3.5. *Let X be a uniformly convex real Banach space and K be a nonempty closed convex subset of X . Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive continuous mappings with sequences $\{\mu_{in}\}_{n=1}^{\infty}, \{\lambda_{in}\}_{n=1}^{\infty}$, where $i = \overline{1, m}$, and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family total asymptotically nonexpansive continuous mappings with sequences $\{\tilde{\mu}_{in}\}_{n=1}^{\infty}, \{\tilde{\lambda}_{in}\}_{n=1}^{\infty}$, where $i = \overline{1, m}$, such that $F := \bigcap_{i=1}^m (F(T_i) \cap F(I_i)) \neq \emptyset$. Suppose $\sum_{n=1}^{\infty} \mu_{in} < \infty$, $\sum_{n=1}^{\infty} \lambda_{in} < \infty$, $\sum_{n=1}^{\infty} \tilde{\mu}_{in} < \infty$, $\sum_{n=1}^{\infty} \tilde{\lambda}_{in} < \infty$ for all $i = \overline{1, m}$ and there exist $M_i, M_i^*, N_i, N_i^* > 0$, $i = \overline{1, m}$ such that $\phi_i(\xi_i) \leq M_i^* \xi_i$, for all $\xi_i \geq M_i$ and $\varphi_i(\zeta_i) \leq N_i^* \zeta_i$ for all $\zeta_i \geq N_i$, where $i = \overline{1, m}$. Then the explicit iterative sequence $\{x_n\}$ defined by (1.15) converges strongly to a common fixed point in F if and only if*

$$\liminf_{n \rightarrow \infty} d(x_n, F) = 0. \quad (3.11)$$

Proof. The necessity of condition (3.11) is obvious. Let us proof the sufficiency part of Theorem.

Since $\{T_i\}_{i=1}^m, \{I_i\}_{i=1}^m : K \rightarrow K$ are continuous mappings, the sets $F(T_i)$ and $F(I_i)$ are closed. Hence $F = \bigcap_{i=1}^m (F(T_i) \cap F(I_i))$ is a nonempty closed set.

For any given $p \in F$, we have (see (3.9))

$$\|x_{n+1} - p\| \leq (1 + b_n) \|x_n - p\| + c_n, \quad (3.12)$$

Hence, one finds

$$d(x_{n+1}, F) \leq (1 + b_n) d(x_n, F) + c_n \quad (3.13)$$

From (3.13) due to Lemma 2.1 we obtain the existence of the limit $\lim_{n \rightarrow \infty} d(x_n, F)$. By condition (3.11), one gets

$$\lim_{n \rightarrow \infty} d(x_n, F) = \liminf_{n \rightarrow \infty} d(x_n, F) = 0.$$

Let us prove that the sequence $\{x_n\}$ converges strongly to a common fixed point in F . We first show that $\{x_n\}$ is Cauchy sequence in X . In fact, due to $1 + t \leq \exp(t)$ for all $t > 0$, and from (3.12), we obtain

$$\|x_{n+1} - p\| \leq \exp(b_n)(\|x_n - p\| + c_n). \quad (3.14)$$

Thus, for any positive integers m, n , from (3.14) with $\sum_{n=1}^{\infty} b_n < \infty$, $\sum_{n=1}^{\infty} c_n < \infty$, we find

$$\begin{aligned} \|x_{n+m} - p\| &\leq \exp(b_{n+m-1})(\|x_{n+m-1} - p\| + c_{n+m-1}) \\ &\leq \exp(b_{n+m-1} + b_{n+m-2})(\|x_{n+m-2} - p\| + c_{n+m-1} + c_{n+m-2}) \\ &\leq \dots \\ &\leq \exp\left(\sum_{i=n}^{n+m-1} b_i\right) \left(\|x_n - p\| + \sum_{i=n}^{n+m-1} c_i\right) \\ &\leq \exp\left(\sum_{i=n}^{\infty} b_i\right) \left(\|x_n - p\| + \sum_{i=n}^{\infty} c_i\right). \end{aligned}$$

Therefore we get

$$\begin{aligned} \|x_{n+m} - x_n\| &\leq \|x_{n+m} - p\| + \|x_n - p\| \\ &\leq \left(1 + \exp\left(\sum_{i=n}^{\infty} b_i\right)\right) \|x_n - p\| + \exp\left(\sum_{i=n}^{\infty} b_i\right) \sum_{i=n}^{\infty} c_i \\ &\leq W \left(\|x_n - p\| + \sum_{i=n}^{\infty} c_i\right) \end{aligned} \quad (3.15)$$

for all $p \in F$, where $0 < W - 1 = \exp\left(\sum_{i=n}^{\infty} b_i\right) < \infty$. Taking infimum over $p \in F$ in (3.15) gives

$$\|x_{n+m} - x_n\| \leq W \left(d(x_n, F) + \sum_{i=n}^{\infty} c_i\right) \quad (3.16)$$

Since $\lim_{n \rightarrow \infty} d(x_n, F) = 0$, and $\sum_{i=1}^{\infty} c_i < \infty$, given $\varepsilon > 0$ there exists an integer $N_0 > 0$ such

that for all $n > N_0$ we have $d(x_n, F) < \frac{\varepsilon}{2W}$ and $\sum_{i=n}^{\infty} c_i < \frac{\varepsilon}{2W}$. Consequently, for all integers $n \geq N_0$ and $m \geq 1$ and from (3.16) we derive

$$\|x_{n+m} - x_n\| \leq \varepsilon.$$

which means that $\{x_n\}$ is Cauchy sequence in X , and since X is complete there exists $x^* \in X$ such that the sequence $\{x_n\}$ converges strongly to x^* .

Now we show that x^* is a common fixed point in F . Suppose for contradiction that $x^* \notin F$. Since F is closed subset of X , we have that $d(x^*, F) > 0$. However, for all $p \in F$, we have

$$\|x^* - p\| \leq \|x_n - x^*\| + \|x_n - p\|.$$

This implies that

$$d(x^*, F) \leq \|x_n - x^*\| + d(x_n, F),$$

so that as $n \rightarrow \infty$ we obtain $d(x^*, F) = 0$ which contradicts $d(x^*, F) > 0$. Hence, x^* is a common fixed point in F . This proves the required assertion. \square

To formulate and prove the main result, we need one more an auxiliary result.

Proposition 3.6. *Let X be a uniformly convex real Banach space and K be a nonempty closed convex subset of X . Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive continuous mappings with sequences $\{\mu_{in}\}_{n=1}^{\infty}, \{\lambda_{in}\}_{n=1}^{\infty}$, where $i = \overline{1, m}$, and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family total asymptotically nonexpansive continuous mappings with sequences $\{\tilde{\mu}_{in}\}_{n=1}^{\infty}, \{\tilde{\lambda}_{in}\}_{n=1}^{\infty}$, where $i = \overline{1, m}$, such that $F := \bigcap_{i=1}^m (F(T_i) \cap F(I_i)) \neq \emptyset$. Suppose $\sum_{n=1}^{\infty} \mu_{in} < \infty, \sum_{n=1}^{\infty} \lambda_{in} < \infty, \sum_{n=1}^{\infty} \tilde{\mu}_{in} < \infty, \sum_{n=1}^{\infty} \tilde{\lambda}_{in} < \infty$ for all $i = \overline{1, m}$, and $\{\alpha_{jn}\}_{n=1}^{\infty}, \{\beta_{jn}\}_{n=1}^{\infty}$ are sequences with $\{\alpha_{jn}\}_{n=1}^{\infty} \subset [\alpha_*, \alpha^*]$ and $\{\beta_{jn}\}_{n=1}^{\infty} \subset [\beta_*, \beta^*]$, for all $j = \overline{0, m}$, here $0 < \alpha_* < \alpha^* < 1, 0 < \beta_* < \beta^* < 1$, and there exist $M_i, M_i^*, N_i, N_i^* > 0, i = \overline{1, m}$ such that $\phi_i(\xi_i) \leq M_i^* \xi_i$, for all $\xi_i \geq M_i$ and $\varphi_i(\zeta_i) \leq N_i^* \zeta_i$ for all $\zeta_i \geq N_i$, where $i = \overline{1, m}$. Then the explicit iterative sequence $\{x_n\}$ defined by (1.15) satisfies the following*

$$\lim_{n \rightarrow \infty} \|x_n - T_i^n x_n\| = 0, \quad (3.17)$$

$$\lim_{n \rightarrow \infty} \|x_n - I_i^n x_n\| = 0, \quad (3.18)$$

for all $i = \overline{1, m}$.

Proof. According to Lemma 3.4 for any $p \in F$ we have $\lim_{n \rightarrow \infty} \|x_n - p\| = d$. It follows from (1.15) that

$$\|x_{n+1} - p\| = \left\| \alpha_{0n}(x_n - p) + \sum_{i=1}^m \alpha_{in}(T_i^n y_n - p) \right\| \rightarrow d, \quad (3.19)$$

as $n \rightarrow \infty$. By means of $\sum_{n=1}^{\infty} \mu_{in} < \infty$, $\sum_{n=1}^{\infty} \lambda_{in} < \infty$, $\sum_{n=1}^{\infty} \tilde{\mu}_{in} < \infty$, $\sum_{n=1}^{\infty} \tilde{\lambda}_{in} < \infty$, for all $i = \overline{1, m}$, from (3.8) one yields that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \|y_n - p\| &\leq \limsup_{n \rightarrow \infty} \left[\left(1 + \sum_{i=1}^m \tilde{\mu}_{in} \beta_{in} N_i^* \right) \|x_n - p\| \right] \\ &\quad + \limsup_{n \rightarrow \infty} \left[\sum_{i=1}^m \left(\tilde{\mu}_{in} \beta_{in} \varphi_i(N_i) + \tilde{\lambda}_{in} \beta_{in} \right) \right] \\ &= \limsup_{n \rightarrow \infty} \|x_n - p\| = d \end{aligned} \quad (3.20)$$

and from (3.4), (3.20) we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \|T_i^n y_n - p\| &\leq \limsup_{n \rightarrow \infty} [(1 + \mu_{in} M_i^*)(1 + \tilde{\mu}_{in} N_i^*) \|y_n - p\| \\ &\quad + \limsup_{n \rightarrow \infty} \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i)] \\ &\quad + \limsup_{n \rightarrow \infty} [\tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) + \mu_{in} \phi_i(M_i) + \lambda_{in}] \\ &\leq d \end{aligned} \quad (3.21)$$

for all $i = \overline{1, m}$. Now using

$$\limsup_{n \rightarrow \infty} \|x_n - p\| = d$$

with (3.21) and applying Lemma 3.2 to (3.19) one finds

$$\lim_{n \rightarrow \infty} \|x_n - T_i^n y_n\| = 0. \quad (3.22)$$

for all $i = \overline{1, m}$. Now from (1.15) and (3.22) we infer that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} \left\| \sum_{i=1}^m \alpha_{in} (T_i^n y_n - x_n) \right\| = 0. \quad (3.23)$$

On the other hand, from (3.4) we have

$$\begin{aligned} \|x_n - p\| &\leq \|x_n - T_i^n y_n\| + \|T_i^n y_n - p\| \\ &\leq \|x_n - T_i^n y_n\| + (1 + \mu_{in} M_i^*)(1 + \tilde{\mu}_{in} N_i^*) \|y_n - p\| \\ &\quad + \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) + \tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) + \mu_{in} \phi_i(M_i) + \lambda_{in} \end{aligned}$$

which implies

$$\begin{aligned} \|x_n - p\| - \|x_n - T_i^n y_n\| &\leq (1 + \mu_{in} M_i^*)(1 + \tilde{\mu}_{in} N_i^*) \|y_n - p\| \\ &\quad + \tilde{\mu}_{in} (1 + \mu_{in} M_i^*) \varphi_i(N_i) + \tilde{\lambda}_{in} (1 + \mu_{in} M_i^*) \\ &\quad + \mu_{in} \phi_i(M_i) + \lambda_{in} \end{aligned}$$

The last inequality with (3.10), (3.22) yields

$$\liminf_{n \rightarrow \infty} \|y_n - p\| = d \quad (3.24)$$

Combining (3.24) with (3.20) we get

$$\lim_{n \rightarrow \infty} \|y_n - p\| = d \quad (3.25)$$

Again from (1.15) we can see that

$$\|y_n - p\| = \left\| \beta_{0n}(x_n - p) + \sum_{i=1}^m \beta_{in}(I_i^n x_n - p) \right\| \rightarrow d, \quad n \rightarrow \infty. \quad (3.26)$$

From (3.3) and (3.10) one finds

$$\limsup_{n \rightarrow \infty} \|I_i^n x_n - p\| \leq \limsup_{n \rightarrow \infty} ((1 + \tilde{\mu}_{in} N_i^*) \|x_n - p\| + \tilde{\mu}_{in} \varphi_i(N_i) + \tilde{\lambda}_{in}) = d$$

for all $i = \overline{1, m}$. Now applying Lemma 3.2 to (3.26) we obtain

$$\lim_{n \rightarrow \infty} \|x_n - I_i^n x_n\| = 0 \quad (3.27)$$

for all $i = \overline{1, m}$. We then have

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = \lim_{n \rightarrow \infty} \left\| \sum_{i=1}^m \beta_{in}(I_i^n x_n - x_n) \right\| = 0 \quad (3.28)$$

Consider

$$\begin{aligned} \|x_n - T_i^n x_n\| &\leq \|x_n - T_i^n y_n\| + \|T_i^n y_n - T_i^n x_n\| \\ &\leq \|x_n - T_i^n y_n\| + (1 + \mu_{in} M_i^*)(1 + \tilde{\mu}_{in} N_i^*) \|y_n - x_n\| \\ &\quad + (1 + \mu_{in} M_i^*)(\tilde{\mu}_{in} \varphi_i(N_i) + \tilde{\lambda}_{in}) + \mu_{in} \phi_i(M_i) + \lambda_{in}, \end{aligned}$$

for all $i = \overline{1, m}$. Then from (3.22) and (3.28) we get

$$\lim_{n \rightarrow \infty} \|x_n - T_i^n x_n\| = 0,$$

for all $i = \overline{1, m}$. □

Now we are ready to formulate a main result concerning strong convergence of the sequence $\{x_n\}$.

Theorem 3.7. *Let X be a uniformly convex real Banach space and K be a nonempty closed convex subset of X . Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically I_i -nonexpansive continuous mappings with sequences $\{\mu_{in}\}_{n=1}^\infty, \{\lambda_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$, and $\{I_i\}_{i=1}^m : K \rightarrow K$ be a finite family total asymptotically nonexpansive continuous mappings with sequences $\{\tilde{\mu}_{in}\}_{n=1}^\infty, \{\tilde{\lambda}_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$, such that $F := \bigcap_{i=1}^m (F(T_i) \cap F(I_i)) \neq \emptyset$. Suppose $\sum_{n=1}^\infty \mu_{in} < \infty$, $\sum_{n=1}^\infty \lambda_{in} < \infty$, $\sum_{n=1}^\infty \tilde{\mu}_{in} < \infty$, $\sum_{n=1}^\infty \tilde{\lambda}_{in} < \infty$ for all $i = \overline{1, m}$, and $\{\alpha_{jn}\}_{n=1}^\infty, \{\beta_{jn}\}_{n=1}^\infty$ are sequences with $\{\alpha_{jn}\}_{n=1}^\infty \subset [\alpha_*, \alpha^*]$ and $\{\beta_{jn}\}_{n=1}^\infty \subset [\beta_*, \beta^*]$, for all $j = \overline{0, m}$, here $0 < \alpha_* < \alpha^* < 1$, $0 < \beta_* < \beta^* < 1$, and there exist $M_i, M_i^*, N_i, N_i^* > 0$, $i = \overline{1, m}$ such that $\phi_i(\xi_i) \leq M_i^* \xi_i$, for all $\xi_i \geq M_i$ and $\varphi_i(\zeta_i) \leq N_i^* \zeta_i$ for all $\zeta_i \geq N_i$, where $i = \overline{1, m}$. If at least one mapping of the mappings $\{T_i\}_{i=1}^m$ and $\{I_i\}_{i=1}^m$ is compact, then the explicitly iterative sequence $\{x_n\}$ defined by (1.15) converges strongly to a common fixed point of $\{T_i\}_{i=1}^m$ and $\{I_i\}_{i=1}^m$.*

Proof. Without any loss of generality, we may assume that T_1 is compact. This means that there exists a subsequence $\{T_1^{n_k}x_{n_k}\}_{k=1}^\infty$ of $\{T_1^n x_n\}_{n=1}^\infty$ such that $\{T_1^{n_k}x_{n_k}\}_{k=1}^\infty$ converges strongly to $x^* \in K$. Then from (3.17) we have that $\{x_{n_k}\}_{k=1}^\infty$ converges strongly to x^* . Also from (3.17), we obtain $\{T_i^{n_k}x_{n_k}\}_{k=1}^\infty$ converges strongly to x^* , for all $i = \overline{2, m}$. Since $\{T_i\}_{i=1}^m$ are continuous mappings, so $\{T_i^{n_k+1}x_{n_k}\}_{k=1}^\infty$ converges strongly to $T_i x^*$, for all $i = \overline{1, m}$. On the other hand, from (3.18) and continuousness of $\{I_i\}_{i=1}^m$ we obtain that $\{I_i^{n_k}x_{n_k}\}_{k=1}^\infty$ converges strongly to x^* and $\{I_i^{n_k+1}x_{n_k}\}_{k=1}^\infty$ converges strongly to $I_i x^*$, for all $i = \overline{1, m}$. Due to (3.23), $\{\|x_{n_k+1} - x_{n_k}\|\}$ converges to 0, as $k \rightarrow \infty$. Then, $\{x_{n_k+1}\}_{k=1}^\infty$ converges strongly to x^* and moreover, (3.4) and (3.3) imply that $\{\|T_i^{n_k+1}x_{n_k+1} - T_i^{n_k+1}x_{n_k}\|\}$ and $\{\|I_i^{n_k+1}x_{n_k+1} - I_i^{n_k+1}x_{n_k}\|\}$ converge to 0, as $k \rightarrow \infty$, for all $i = \overline{1, m}$. From (3.17), (3.18) it yields that $\|x_{n_k+1} - T_i^{n_k+1}x_{n_k+1}\|$ and $\|x_{n_k+1} - I_i^{n_k+1}x_{n_k+1}\|$ converge to 0 as $k \rightarrow \infty$, for all $i = \overline{1, m}$. Observe that

$$\begin{aligned} \|x^* - T_i x^*\| &\leq \|x^* - x_{n_k+1}\| + \|x_{n_k+1} - T_i^{n_k+1}x_{n_k+1}\| \\ &\quad + \|T_i^{n_k+1}x_{n_k+1} - T_i^{n_k+1}x_{n_k}\| + \|T_i^{n_k+1}x_{n_k} - T_i x^*\|, \\ \|x^* - I_i x^*\| &\leq \|x^* - x_{n_k+1}\| + \|x_{n_k+1} - I_i^{n_k+1}x_{n_k+1}\| \\ &\quad + \|I_i^{n_k+1}x_{n_k+1} - I_i^{n_k+1}x_{n_k}\| + \|I_i^{n_k+1}x_{n_k} - I_i x^*\|, \end{aligned}$$

for all $i = \overline{1, m}$. Taking limit as $k \rightarrow \infty$ we have that $x^* = T_i x^*$ and $x^* = I_i x^*$, for all $i = \overline{1, m}$, which means $x^* \in F$. However, by Lemma 3.4, the limit $\lim_{n \rightarrow \infty} \|x_n - x^*\|$ exists then

$$\lim_{n \rightarrow \infty} \|x_n - x^*\| = \lim_{n_k \rightarrow \infty} \|x_{n_k} - x^*\| = 0,$$

which means $\{x_n\}$ converges strongly to $x^* \in F$. This completes the proof. \square

Remark. If one has that all I_i are identity mappings, then the obtained results recover and correctly prove the main result of [11].

Remark. Suppose we are given two family $\{T_i\}_{i=1}^m : K \rightarrow K$ and $\{S_i\}_{i=1}^m : K \rightarrow K$ of total asymptotically nonexpansive continuous mappings such that $\bigcap_{i=1}^m (F(T_i) \cap F(S_i)) \neq \emptyset$. Define the following explicit iterative process:

$$\begin{cases} x_0 \in K, \\ x_{n+1} = \alpha_{0n}x_n + \sum_{i=1}^m \alpha_{in}T_i^{n_n}y_n + \alpha_{m+1,n}u_n \\ y_n = \beta_{0n}x_n + \sum_{i=1}^m \beta_{in}S_i^{n_n}x_n + \beta_{m+1,n}v_n. \end{cases} \quad (3.29)$$

such that $\sum_{j=0}^{m+1} \alpha_{jn} = 1$ and $\sum_{j=0}^{m+1} \beta_{jn} = 1$.

Under suitable conditions, by the same argument and methods used above one can prove, with either little mirror or no modifications, the strong convergence of the explicit iterative process defined by (3.29) to a common fixed point of the given families.

Remark. Let $\{T_i\}_{i=1}^m : K \rightarrow K$ be a finite family of total asymptotically nonexpansive continuous mappings with sequences $\{\mu_{in}\}_{n=1}^\infty, \{\lambda_{in}\}_{n=1}^\infty$, where $i = \overline{1, m}$. It is clear for each operator T_i one has

$$\|T_i^n x - T_i^n y\| \leq \|T_i^n x - T_i^n y\| + \mu_{in} \|T_i^n x - T_i^n y\|, \quad (3.30)$$

this means that T_i is total asymptotically T_i -nonexpansive mappings with sequence $\{\mu_{in}\}_{n=1}^\infty$ and the function $\phi(\lambda) = \lambda$. Hence, our iteration scheme can be written as follows

$$\begin{cases} x_0 \in K, \\ x_{n+1} = \alpha_{0n}x_n + \sum_{i=1}^m \alpha_{in}T_i^n y_n \\ y_n = \beta_{0n}x_n + \sum_{i=1}^m \beta_{in}T_i^n x_n. \end{cases} \quad (3.31)$$

where $\{\alpha_{jn}\}_{n=1}^\infty, \{\beta_{jn}\}_{n=1}^\infty$ in $(0, 1)$, $(j = \overline{0, m})$ with $\sum_{j=0}^m \alpha_{jn} = 1, \sum_{j=0}^m \beta_{jn} = 1$.

The defined scheme is a new iterative method generalizing one given in [11]. So, according our main results for the defined sequence $\{x_n\}$ (see (3.31)) we obtain strong convergence theorems. On the other hand, playing with numbers $\{\alpha_{jn}\}_{n=1}^\infty, \{\beta_{jn}\}_{n=1}^\infty$ and by means of the defined method one may introduce lots of different schemes. All of the them strongly converges to a common fixed point of $\{T_i\}_{i=1}^m$. Moreover, the recursion formula (3.31) is much simpler than the others studied earlier for this problem [1], [10], [18], [35],[28],[2],[24],[19],[40]. Therefore, all presented results here generalize, unify and extend the corresponding main results of the mentioned papers. Note that one can consider the method (1.15) with errors, and all the theorems could be carry over for such iteration scheme as well with little or no modifications.

We stress that all the theorems of this paper carry over to the class of total asymptotically quasi I -nonexpansive mappings (see [11]), [39] with little or no modifications.

Acknowledgments

A part of this work was done at the Abdus Salam International Center for Theoretical Physics (ICTP), Trieste, Italy. The first named author (FM) thanks the ICTP for providing financial support during his visit as a Junior Associate at the centre. The authors also acknowledge the Malaysian Ministry of Science, Technology and Innovation Grant 01-01-08-SF0079.

References

- [1] Ya.I. Alber, C.E. Chidume, H. Zegeye, Approximating fixed points of total asymptotically nonexpansive mappings, *Fixed Point Theory Appl.* **2006** (2006), Article ID 10673, 20 p.

- [2] Ya.I. Alber, R. Espinola, P. Lorenzoc, Strongly Convergent Approximations to Fixed Points of Total Asymptotically Nonexpansive Mappings, *Acta Math. Sinica, Eng. Ser.* **24** (2008) 1005-1022.
- [3] F.E. Browder, Nonexpansive nonlinear operators in Banach space, *Proc. Natl. Acad. Sci. USA* **54** (1965) 1041–1044.
- [4] F.E. Browder, Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach space, *Arch. Ration. Mech. Anal.* **24** (1967) 82–90.
- [5] R.E. Bruck, Asymptotic behavior of nonexpansive mappings, *Contemporary Mathematics, 18, Fixed Points and Nonexpansive Mappings*, AMS, Providence, RI, 1980.
- [6] B. Bruck., T. Kuczumow, S. Reich, Convergence of iterates of asymptotically nonexpansive mappings in Banach space with the uniform Opial property, *Colloq. Math.* **65** (1993) 169–179.
- [7] S. S. Chang, R. P. Agarwal, Y. J. Cho, Strong convergence of iterative sequences for asymptotically nonexpansive mappings in Banach spaces, *Dynamic Systems Appl.* **14**(2005), 447–454.
- [8] S. S. Chang, Y. J. Cho, Y. X. Tian, Strong convergence theorems of Reich type iterative sequence for non-self asymptotically nonexpansive mappings, *Taiwan. J. Math.* **11**(2007), 729–743.
- [9] C.E. Chidumi, *Geometric Properties of Banach Spaces and Nonlinear Iterations*, Lecture Note Math. Vol. 1965, Springer-Verlag London, 2009.
- [10] C.E. Chidume, E.U. Ofoedu, Approximation of common fixed points for finite families of total asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **333** (2007) 128–141.
- [11] C.E. Chidume, E.U. Ofoedu, A new iteration process for approximation of common fixed points for finite families of total asymptotically nonexpansive mappings, *Inter. J. Math. & Math. Sci.* **2009** (2009) Article ID 615107, 17p.
- [12] Y. J. Cho, N. Hussain, H. K. Pathak, Approximation of nearest common fixed points of asymptotically I -nonexpansive mappings in Banach spaces, *Commun. Korean Math. Soc.* **26**(2011), 483–498.
- [13] Y. J. Cho, J. I. Kang, H. Y. Zhou, Approximating common fixed points of asymptotically nonexpansive mappings, *Bull. Korean Math. Soc.* **42**(2005), 661-670.
- [14] Y. J. Cho, H. Y. Zhou, G. T. Guo, Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings, *Comput. Math. Appl.* **47**(2004), 707–717.

- [15] K. Goebel, W.A. Kirk, *Topics in metric fixed point theory*, Cambridge Studies in Advanced Mathematics 28, CUP, Cambridge, New York, Port Chester, Melbourne, Sydney, 1990.
- [16] K. Goebel, W.A. Kirk, A fixed point theorem for asymptotically nonexpansive mapping, *Proc. Amer. Math. Soc.* **35** (1972) 171–174.
- [17] J. Gornicki, Weak convergence theorems for asymptotically nonexpansive mappings in uniformly Banach space, *Comment. Math. Univ. Carolinae* **30** (1989) 249–252.
- [18] F. Gu, Some convergence theorems of non-implicit iteration process with errors for a finite families of I -asymptotically nonexpansive mappings, *Appl. Math. Comp.* **216** (2010), 161–172.
- [19] W. Guo, Y. J. Cho, On the strong convergence of the implicit iterative processes with errors for a finite family of asymptotically nonexpansive mappings, *Appl. Math. Lett.* **21**(2008), 1046–1052.
- [20] W.A. Kirk W.A., Fixed point theorems for non-Lipschitzian mappings of asymptotically nonexpansive type, *Israel J. Math.* **17** (1974) 339–346.
- [21] G.E. Kim, T.H. Kim, Mann and Ishikawa iterations with errors for non-Lipschitzian mappings an Banach space, *Computers Math. Appl.* **42** (2001) 1565–1570.
- [22] Q. Liu, Iterative sequence for asymptotically quasi-nonexpansive mappings, *J. Math. Anal. Appl.* **256** (2001) 1-7.
- [23] P. Kumam, W. Kumethong, N. Jewwaiworn, Weak convergence theorems of three-step Noor iterative scheme for I -quasi-nonexpansive mappings in Banach spaces, *Appl. Math. Sci.*, **2** (2008) 2915– 2920.
- [24] F.Mukhamedov, M.Saburov, Weak and strong convergence of an implicit iteration process for an asypmtotically quasi- I -nonexpansive mapping in Banach space, *Fixed Point Theory Appl.* **2010**(2010), Article ID 719631, 13 p.
- [25] F.Mukhamedov, M.Saburov, Strong convergence of an explicit iterative process for a totally asymptotically I -nonexpansive mappings in Banach spaces, *Appl. Math. Letters* **23** (2010), 1473–1478.
- [26] S. Reich, Strong convergence theorems for resolvents of accretive operators in Banach spaces, *J. Math. Anal. Appl.* **75** (1980) 287-292.
- [27] B.H. Rhodes, S. Temir, Convergebce thorems for I -nonexpansive mapping, *Inter. Jour. Math. & Math. Sci.* **2006** (2006), Article ID 63435, 1-4.

- [28] D.R. Sahu, Fixed points of demicontinuous nearly Lipschitzian mappings in Banach spaces, *Commen. Math. Univ. Carolinae* **46** (2005) 653–666.
- [29] N. Shahzad, A. Udomene, Approximating common fixed points of two asymptotically quasi-nonexpansive mappings in Banach spaces, *Fixed Point Theory Appl.* **2006** (2006) Article ID 18909, 10p.
- [30] N. Shahzad, Generalized I -nonexpansivemaps and best approximations in Banach spaces, *Demonstratio Math.* **37** (2004) 597-600.
- [31] J. Schu, Weak and strong convergence to fixed points of asypmtotically nonexpansive mappings, *Bull. Australian Math. Soc.* **43** (1991) 153–159.
- [32] N. Shoji, W. Takahashi, Strong convergence theorems for asymptotically nonexpansive semigroups in Hilbert spaces, *Nonlinear Anal.* **34** (1998) 87-99.
- [33] K.-K. Tan, H.-K. Xu, Approximating fixed points of nonexpansive mapping by the Ishikawa iteration process, *J. Math. Anal. Appl.* **178** (1993) 301–308.
- [34] S. Temir, O. Gul, Convergence theorem for I -asymptotically quasi-nonexpansive mapping in Hilbert space, *J. Math. Anal. Appl.* **329** (2007) 759–765.
- [35] S. Temir, On the convergence theorems of implicit iteration process for a finite family of I -asymptotically nonexpansive mappings, *J. Comp. Appl. Math.* **225** (2009) 398–405.
- [36] S. Temir, Convergence of Iterative Process for Generalized I - asymptotically Quasi-Nonexpansive Mappings *Thai J. Math* **7** (2009) 367-379.
- [37] R. Wittmann, Approximation of the fixed points of nonexpansive mappings, *Arch. Math.* **58** (1992) 486–491.
- [38] E. Zeidler, *Nonlinear Functional Analysis and Its Applications. I: Fixed-Point Theorems*, Springer, New York, NY, USA, 1986.
- [39] J.Zhao, S. He, G. Liu, Strong convergence theorems for generalized asymptotically quasi-nonexpansive mappings, *J Appl Math Comput* **30** (2009) 53-64.
- [40] H. Y. Zhou, Y. J. Cho, S. M. Kang, A new iterative algorithm for approximating common fixed points for asymptotically nonexpansive mappings, *Fixed Point Theory Appl.* Vol. 2007, Article ID 64974, pp. 10.